CONTENTS

	Page
Fusarium-Blight (Scab) of Wheat and Other Cereals. DIMITR ATANASOFF	1
Cause of Lime-Induced Chlorosis and Availability of Iron in the Soil. P. L. GILE and J. O. CARRERO	33
An Experimental Study of Echinacea Therapy. JAMES F. COUCH and LEIGH T. GILTNER	63
Investigations of the Germicidal Value of Some of the Chlorin Disinfectants. F. W. TILLEY	85
A New Avocado Weevil from the Canal Zone. H. F. DIETZ and H. S. BARBER	111
Studies in Mustard Seeds and Substitutes: I. Chinese Colza (Brassica campestris chinoleifera Viehoever). ARNO VIEHOEVER, JOSEPH F. CLEVENGER, and CLARE OLIN EWING	117
Study of Some Poultry Feed Mixtures with Reference to Their Potential Acidity and Their Potential Alkalinity. B. F. KAUPP and J. E. IVEY	141
The Influence of Cold in Stimulating the Growth of Plants. FREDERICK V. COVILLE	151
Composition of Normal and Mottled Citrus Leaves. W. P. KELLEY and A. B. CUMMINS	161
Control of Fluke Diseases by Destruction of the Intermediate Host. ASA C. CHANDLER	193
Injury to Seed Wheat Resulting from Drying after Disinfection with Formaldehyde. ANNIE MAY HURD	209
Studies on the Life History and Habits of the Beet Leafhopper. C. F. STAHL	245
Hypertrophied Lenticels on the Roots of Conifers and Their Relation to Moisture and Aeration. GLENN G. HAHN, CARL HARTLEY, and ARTHUR S. RHOADS	253
Degree of Temperature to Which Soils Can Be Cooled without Freezing. GEORGE BOUYOUCOS	267
Changes Taking Place in the Tempering of Wheat. E. L. TAGUE	271
Vascular Discoloration of Irish Potato Tubers. H. A. EDSON	277
Crownwart of Alfalfa Caused by Urophlyctis alfalfae. FRED RUEL JONES and CHARLES DRECHSLER	295
Pathological Anatomy of Potato Blackleg. ERNST F. ARTSCHWAGER	325

Page

Sclerotinia minor, n. sp., the Cause of a Decay of Lettuce, Celery, and Other Crops. IVAN C. JAGGER 331

Permanence of Differences in the Plots of an Experimental Field. J. ARTHUR HARRIS and C. S. SCOFIELD 335

Some Changes in Florida Grapefruit in Storage. LON A. HAWKINS and J. R. MAGNESS 357

A Bacteriological Study of Canned Ripe Olives. STEWART A. KOSER 375

Relation of the Soil Solution to the Soil Extract. D. R. HOAGLAND, J. C. MARTIN, and G. R. STEWART 381

Effect of Season and Crop Growth on the Physical State of the Soil. D. R. HOAGLAND and J. C. MARTIN 397

Carbon-Dioxid Content of Barn Air. MARY F. HENDRY and ALICE JOHNSON 405

Rice Weevil, (Calandra) Sitophilus oryza. RICHARD T. COTTON. 409

Opius fletcheri as a Parasite of the Melon Fly in Hawaii. H. F. WILLARD 423

Tamarind Pod-Borer, Sitophilus linearis (Herbst). RICHARD T. COTTON 439

Influence of Temperature and Humidity on the Growth of Pseudomonas citri and Its Host Plants and on Infection and Development of the Disease. GEORGE L. PELTIER 447

Daubentonia longifolia (Coffee Bean), A Poisonous Plant. C. DWIGHT MARSH and A. B. CLAWSON 507

Fusarium-Wilt of Tobacco. JAMES JOHNSON 515

Sugar Beet Top Silage. RAY E. NEIDIG 537

Nodule Bacteria of Leguminous Plants. F. LÖHNIS and ROY HANSEN 543

Correlation and Causation. SEWALL WRIGHT 557

Measurement of the Amount of Water That Seeds Cause to Become Unfree and Their Water-Soluble Material. GEORGE J. BOUYOUCOS and M. M. MCCOOL 587

Inheritance of Syndactylism, Black, and Dilution in Swine. J. A. DETLEFSEN and W. J. CARMICHAEL 595

Four Rhynchophora Attacking Corn in Storage. RICHARD T. COTTON 605

Concentration of Potassium in Orthoclase Solutions Not a Measure of Its Availability to Wheat Seedlings. J. F. BREAZEALE and LYMAN J. BRIGGS 615

Composition of Tubers, Skins, and Sprouts of Three Varieties of Potatoes. F. C. COOK 623

Further Studies in the Deterioration of Sugars in Storage. NICHOLAS KOPELOFF, H. Z. E. PERKINS, and C. J. WELCOME 637

Freezing of Fruit Buds. FRANK L. WEST and N. E. EDLEFSEN .. 655

Page
Effect of Various Crops Upon the Water Extract of a Typical Silty Clay Loam Soil. G. R. STEWART and J. C. MARTIN 663
Another Conidial Sclerospora of Philippine Maize. WILLIAM H. WESTON, Jr 669
Onion Smudge. J. C. WALKER 685
Variations in Colletotrichum gloeosporioides. O. F. BURGER 723
A Transmissible Mosaic Disease of Lettuce. IVAN C. JAGGER .. 737
Leconte's Sawfly, an Enemy of Young Pines. WILLIAM MIDDLETON 741
Amylase of Rhizopus tritici, with a Consideration of Its Secretion and Action. L. L. HARTER 761
A Comparative Study of the Composition of the Sunflower and Corn Plants at Different Stages of Growth. R. H. SHAW and P. A. WRIGHT 787
Evaluation of Climatic Temperature Efficiency for the Ripening Processes in Sweetcorn. CHARLES O. APPLEMAN and S. V. EATON 795
Some Lepidoptera Likely to Be Confused with the Pink Bollworm. CARL HEINRICH 807
Biology of the Smartweed Borer, Pyrausta ainsliei Heinrich. GEORGE G. AINSLIE and W. B. CARTWRIGHT 837
Effect of X-Rays on Trichinæ. BENJAMIN SCHWARTZ 845
Relation of the Calcium Content of Some Kansas Soils to the Soil Reaction as Determined by the Electrometric Titration. C. O. SWANSON, W. L. LATSHAW, and E. L. TAGUE 855
Green Feed versus Antiseptics as a Preventive of Intestinal Disorders of Growing Chicks. A. G. PHILIPS, R. H. CARR, and D. C. KENNARD 869
Comparative Utilization of the Mineral Constituents in the Cotyledons of Bean Seedlings Grown in Soil and in Distilled Water. G. DAVIS BUCKNER 875
Sunflower Silage Digestion Experiment with Cattle and Sheep. RAY E. NEIDIG, C. W. HICKMAN, and ROBERT S. SNYDER 881
Index 889

ERRATA AND AUTHORS' EMENDATIONS

Page 9, line 23, "spikelet" should read "group."
Page 16, line 36, "*Triticum*" should read "*Agropyron.*"
Page 68, after line 22 insert "picture."
Page 84, last line, omit "Not seen."
Page 124, line 23, omit "15."
Page 140, legend of Plate 15, omit "Late rosette stage of Chinese colza seedling."
Pages 166, 168, 170, 175, 178, 180, and 181, Tables II, IV, VI, IX, XI, XIII, and XV, after column heading "ash" insert "expressed as percentages of dry matter."
Pages 183 and 184, Tables XVI, XVII, and XVIII, column 2, omit "per cent."
Page 193, line 16, "epidemic" should read "endemic."
Page 200, line 23, "0.005" should read "0.0005."
Page 236, footnote to Table XII, after "42 days" insert "and 60 days."
Page 246, line 23, after "apical portion" insert "of claval region."
Page 413, Table I, cloumn 10, footnote reference "e" should read "l."
Page 414, Table I, column 6, footnote reference "o" should be transposed to column 7.
Page 422, citation 4, omit "In press" and insert "no. 9, p. 235–243."
Page 452, Table II, line 1, footnote reference "a" should be inserted before all entries.
Page 479, line 32, "organism" should read "organisms."
Page 481, "26°" should read "25°."
Page 491, "Table XVII" should read "Table XVIII."
Page 508, Table I, column 3, lines 13 to 22, ".283" should read ".028."
Page 607, line 13, "molas" should read "molar."
Page 614, Plate 72, figure E, and Plate 74, figure E, "ae" should read "al."
Page 810, line 25, "Kostelezkya" should read "Kosteletzkya."
Page 815, line 32, "divini" should read "diveni."
Page 816, line 18, "hessitans" should read "haesitans."
Page 822, line 38, "Kosteleyzkya" should read "Kosteletzkya."
Page 828, line 32, "Kostelelzkya" should read "Kosteletzkya."

ILLUSTRATIONS

Fusarium-Blight (Scab) of Wheat and Other Cereals

Text Figures

Page

1. Conidia of *Gibberella saubinetii*..... 16
2. Special culture tube for maintaining moisture in culture..... 18

Plates

1. *Gibberella saubinetii:* Blighted ("scabbed") wheat heads..... 32
2. *Gibberella saubinetii:* A.—Footrot of wheat caused by Fusarium. B.—Seedling-blight of wheat caused by *G. saubinetii*..... 32
3. A.—Fusarium seedling-blight. B.—Tissue invaded by *G. saubinetii* in causing the headblight of wheat..... 32
4. *Gibberella saubinetii:* A.—Kernels blighted and shriveled by Fusarium-blight. B.—Perithecia development of *G. saubinetii* on an infected wheat head..... 32

Cause of Lime-Induced Chlorosis and Availability of Iron in the Soil

Plates

5. A.—Rice grown in calcareous and noncalcareous soils and sprayed with ferrous sulphate solution (experiment I). B.—Apparatus used in growing plants in experiment VII..... 62
6. A.—Effect of carbonate of lime in depressing the availability of iron (experiment VII). B.—Effect of various substances on growth of rice in calcareous soil (experiment VIII)..... 62

A New Avocado Weevil from the Canal Zone

Plates

7. *Heilipus perseae:* A, B.—Adult, paratype. C.—An avocado fruit (reduced) showing feeding injury by the beetles..... 116
8. *Heilipus perseae:* Leaves showing the injury done by five beetles in 48 hours..... 116
9. *Heilipus perseae*, mature larva: A.—Ventral face of ventral mouthparts. B.—Anterior part of head from above. C.—Lingua, hypopharynx, hypopharyngeal bracon, and dorsal (buccal) face of maxilla. D.—Dorsal face of mandible. E.—Epipharynx. F.—Ventral face of mandible. G.—Head capsule from above. H.—Thoracic spiracle from outside. I.—Mature larva..... 116

Studies in Mustard Seeds and Substitutes: I. Chinese Colza (Brassica campestris chinoleifera Viehoever)

Plates

10. A.—Yellow seed of Chinese colza. B.—Brown seed of Chinese colza. C.—Surface section of yellow seed of Chinese colza, showing lack of reticulations. D.—Surface section of brown seed of Chinese colza, showing reticulations. E.—Cross section of yellow seed of Chinese colza. F.—Cross section of brown seed of Chinese colza..... 140

Page

11. Seedling of Chinese colza, showing cotyledons and young leaves..... 140
12. Early rosette stage of Chinese colza seedling: A.—Plants from (1) brown seed and (2) yellow seeds. B.—Usual form, showing almost entire leaves..... 140
13. Early rosette stage of Chinese colza seedling: A.—Plant showing a variation in lobing of the leaves. Two months old. B.—Plant showing a variation in lobing of the leaves. Three months old..... 140
14. Late rosette stage of Chinese colza seedling: A.—Usual form. B.—Plant showing a variation in lobing of the leaves..... 140
15. A.—Pe-tsai. B.—Cross between Pak-choi and Pe-tsai. C.—Pak-choi.... 140
16. Early flowering stage of Chinese colza: A.—Usual form, showing somewhat enlarged stem base and stem-clasping leaves. B.—Plant without enlarged stem base. C.—Usual form, showing glaucous leaves..... 140
17. Early flowering stage of Chinese colza: A.—Usual form, showing luxuriant growth and long pedicels. B.—Flower cluster..... 140
18. A.—Fruiting stage of Chinese colza. B.—Mature fruit of Chinese colza... 140
19. A.—Herbarium specimen of *Brassica chinensis* L. B.—Herbarium specimen of *Brassica campestris*..... 140

THE INFLUENCE OF COLD IN STIMULATING THE GROWTH OF PLANTS

Plates

20. A.—Blueberry plants, *Vaccinium corymbosum*, made dormant without cold. B.—Chilled and unchilled blueberry plants..... 160
21. A.—Chilled and unchilled plants of grouseberry, *Viburnum americanum*. B.—Chilled and unchilled plants of tamarack, *Larix laricina*..... 160
22. A.—Chilled and unchilled plants of wild crab, *Malus coronaria*. B.—Blueberry plant with one branch stimulated to growth by cold..... 160
23. Blueberry plant with one branch kept dormant by heat. A.—Dormant indoor blueberry plant as it appeared on February 15, 1912. B.—Same plant photographed May 21..... 160
24. A.—Blueberry cuttings starting to grow at 36° F. B.—Blueberry plant growing in the dark at 36° F..... 160
25. A.—Dormant wild crab stimulated to growth by pruning. B.—Dormant wild crabs stimulated to growth by girdling and by notching the stem.. 160
26. A.—Dormant blueberry buds stimulated to growth by chalking the stem. B.—Dormant blueberry bud stimulated to growth by rubbing the stem. 160
27. A.—Normal spring growth on a blueberry stem. B.—Abnormal spring growth on a blueberry stem, due to lack of chilling..... 160
28. Blueberry leaf exuding sugar from glands interpreted as osmotic-pressure safety valves..... 160
29. A plant of bunchberry, *Cornus canadensis*, the seeds of which do not germinate without chilling..... 160
30. A.—Trailing arbutus, *Epigaea repens*, flowering sparingly from lack of chilling. B.—Trailing arbutus plant flowering normally after chilling. C.—Blueberry plant forced into flower in September by artificial chilling.. 160
31. A.—Abnormal growth of an unchilled blueberry plant. B.—Awakening of long dormant plants by artificial chilling..... 160
32. Plants brought out of dormancy at a specified time. A.—Blueberry plants from a lot that had been kept in a dormant condition by warmth for nearly a year. B.—Representative plants from each of the two chilled lots described under A, from photograph made January 18, 1918..... 160

33. A.—Plantation at Whitesbog, N. J., for the testing of blueberry hybrids. B.—Four-year-old blueberry hybrid in full fruit ..... Page 160
34. The ordinary wild blueberry of New Jersey 160
35. Fruit of a selected hybrid blueberry 160

INJURY TO SEED WHEAT RESULTING FROM DRYING AFTER DISINFECTION WITH FORMALDEHYDE

Text Figures

1. Graph showing rate of evaporation of paraformaldehyde at room temperature, approximately 20° C 222
2. Graph showing the relation of humidity of the air to percentage of germination of stored seed in first experiment 226
3. Graph showing the relation of humidity of the air to percentage of germination of stored seed in second experiment 228
4. Graph showing the relation between humidity of the air and seed injury as indicated by rate of growth of germinated seedlings 229
5. Graph showing the diminution in the rate of evaporation of paraformaldehyde inclosed in a desiccator of 2,400-cc. volume 232

Plates

36. A.—Post-treatment seed injury occurring when wheat is dried after treatment with a 0.1 per cent solution. B.—Germinating seedlings of Little Club wheat, showing characteristic post-treatment injury when seed is treated with a 0.1 per cent solution 244
37. A.—Pots showing germination of treated seed stored for 32 days after disinfection with a 0.1 per cent solution of formaldehyde. B.—Wheat plants grown in soil from seed stored for 60 days after disinfection with a 0.1 per cent solution of formaldehyde 244
38. A.—Wheat seedlings showing injury produced by allowing the seed to lie in dry soil for 30 days after treatment with a 0.1 per cent solution of formaldehyde. B.—Desiccators with different degrees of atmospheric humidity obtained by the use of mixtures of sulphuric acid and water in different proportions 244
39. Germinating samples of wheat stored for 35 days after treatment in the desiccators shown in Plate 38 B, illustrating the relation of seed injury to humidity 224
40. Varying injury to wheat treated with a 0.1 per cent solution of formaldehyde, and stored in sealed bottles. A.—Sealed immediately after treatment, 100 per cent germination. B.—Sealed after drying 7 hours, spread on towels in laboratory, no germination. C.—Sealed after drying 24 hours, spread on towels in laboratory, no germination. D.—Sealed after drying 3 days, spread on towels in laboratory, 14 per cent germination 244
41. Germinating wheat kernels, showing the prevention of post-treatment injury by washing the seed with water immediately after treatment. A.—Treated with 0.2 per cent solution, which was not washed off before drying, 32 per cent germination. B.—Treated with 0.2 per cent solution which was washed off before drying, 76 per cent germination. C.—Treated with 0.1 per cent solution, which was not washed off before drying, 52 per cent germination. D.—Treated with 0.1 per cent solution, which was washed off before drying, 74 per cent germination 244

STUDIES ON THE LIFE HISTORY AND HABITS OF THE BEET LEAFHOPPER

Plates

42. *Eutettix tenella:* A.—Adult, light form. B.—Adult, dark form. C.—Adult, color gradation between A and B. D.—Nymph with protruding sac of dryinid parasite..... Page 252
43. Parasites of *Eutettix tenella:* A.—*Pipunculus industrius:* Adult, much enlarged. B.—*Polynema eutettixi:* Adult, much enlarged..... 252

HYPERTROPHIED LENTICELS ON THE ROOTS OF CONIFERS AND THEIR RELATION TO MOISTURE AND AERATION

Plates

44. Section through a hypertrophied lenticel on root of *Pinus rigida* growing in swampy situation..... 266
45. A.—Hypertrophied lenticels on the basal part of layering stem of *Picea mariana*, which had been covered with sphagnum. B.—Tap root of a *Pinus ponderosa* transplant, bearing an unusually large number of hypertrophied lenticels..... 266
46. A.—Cross section of the stem through one of the hypertrophied lenticels shown in C. B.—Large patches of excrescences upon the tap root near the root crown, on *Pinus rigida*. C.—Hypertrophied lenticels on root of 5-months-old *Pinus ponderosa*, grown in a loosely stoppered 2-ounce bottle, in tap water which had not been changed since the germination of the seed..... 265

CROWNWART OF ALFALFA CAUSED BY UROPHLYCTIS ALFALFAE

Plates

47. *Urophlyctis alfalfae:* Drawing of alfalfa plant, showing abundance of crownwart, as found early in May, 1919, in northern California..... 324
48. *Urophlyctis alfalfae:* A–D.—Peripheral portions of actively growing thallus of parasite dissected from living host. E.—Nearly mature resting spore viewed from distil side, showing 11 haustoria in zonate arrangement. F.—Mature resting spore viewed from distil pole, showing 13 pits that mark former location of haustoria. G.—Mature resting spore viewed in profile, showing pits in zonate arrangement and light concavity on proximal side of spore..... 324
49. *Urophlyctis alfalfae:* A.—Section of epidermal region of young foliar structures, showing young primary turbinate cells *ta-tg*, the first products of infection, within epidermal cells. B.—Section of young foliar element, showing wall of invaded epidermal cell disrupted and advance of secondary turbinate cells *tbc-tbe* into underlying tissue. C.—Section of turbinate cell, showing 3 evacuated peripheral segments *pa-pc*. D.—Section of maturing resting spore, showing 8 nuclei and a central vacuole containing 4 granules staining red. E.—Section of mature resting spore, showing numerous red-staining granules in center and 5 nuclei. F.—Section of maturing resting spore, showing 11 normal nuclei and 4 enlarged nuclei in center, the latter apparently degenerating..... 324
50. *Urophlyctis alfalfae:* Section of diseased bud scale of alfalfa, showing four coalescing cavities, in three of which the large primary turbinate cells *taa*, *tba*, and *tc* may be distinguished..... 324
51. *Urophlyctis alfalfae:* Section of diseased bud scale attacked by *U. alfalfae*, showing a group of eight well-developed cavities *a-h* and their relation to the host tissue..... 324

52. A, C, D.—*Urophlyctis pluriannulatus*. B.—*Urophlyctis alfalfae*. A.—Portion of actively growing thallus of *U. pluriannulatus* dissected from gall on leaf of *Sanicula menziesii*, including a turbinate cell *ta* with a nearly mature resting spore *ra*. B.—Abnormally enlarged hyphae and turbinate cells of *U. alfalfae*, showing conspicuous thickening of the walls. C.—Peripheral portion of actively growing thallus of *U. pluriannulatus*, similar to A, showing 8 turbinate cells of the second order, of which 7 have produced turbinate cells of the last order as well as resting spores. D.—Nearly mature resting spore of *U. pluriannulatus*, viewed from polar end, showing 22 haustoria in zonate arrangement.... Page 324

53. *Urophlyctis pluriannulatis:* Section of leaf of *Sanicula menziesii*, showing development of parasite within gall..... 324

54. Crowns of alfalfa plants bearing galls caused by *Urophlyctis alfalfae* photographed at different stages of development. A.—A comparatively early stage of development at which the origin of the gall structures from the elements of developing buds can be traced. B.—A later stage of development at which the origin of the tissue has become obscured..... 324

55. A comparatively early stage of host reaction to invasion by *Urophlyctis alfalfae*..... 324

56. A.—Late stage of development of host reaction to the invasion of *Urophlyctis alfalfae*. B.—Vertical section through a well-developed gall near its central axis, showing its laminated structure arising from the thickening of bud elements..... 324

PATHOLOGICAL ANATOMY OF POTATO BLACKLEG

Text Figure

1. Section of potato leaf, showing distribution of protein crystals..... 329

Plates

57. A.—Plant affected with blackleg. B.—Section of single upper epidermal cell of leaf and adjacent palisade cell. C.—Section of pith cell which is transformed into a sclereid adjacent to phloem fibers..... 330

58. A.—Pith cells of petiole transformed into sclereids with typically stratified walls. B.—Vascular tissue of the petiole greatly increased by blackleg.. 330

SCLEROTINIA MINOR, N. SP., THE CAUSE OF A DECAY OF LETTUCE, CELERY, AND OTHER CROPS

Text Figure

1. Camera lucida drawings of *S. minor:* A, Microconidia and conodiophores; B, Ascospores; C, Germinating ascospores; D, Asci and paraphyses..... 332

Plate

59. A.—Sclerotia on hard potato agar: center, *Sclerotina libertiana*, either end, *S. minor*. B.—Apothecia of *S. libertiana*. C.—Apothecia of *S. minor*.. 334

RELATION OF THE SOIL SOLUTION TO THE SOIL EXTRACT

Text Figures

1. Graph showing relation of freezing-point depressions in soil (calculated to 22 per cent moisture) to total solids extracted by 5 parts of water to 1 of soil..... 382

2. Graph showing relation of freezing-point depressions in soil (calculated to 17 per cent moisture) to total solids extracted by 5 parts of water to 1 of soil ..... Page 383

EFFECT OF SEASON AND CROP GROWTH ON THE PHYSICAL STATE OF THE SOIL

Text Figures

1. Effect of crop on physical state and electrolyte concentration of the water extract of the soil 398
2. Effect of crop on physical state and electrolyte concentration of the water extract of the soil 399
3. Effect of crop on physical state and electrolyte concentration of the water extract of the soil 400
4. Effect of crop on physical state and electrolyte concentration of the water extract of the soil 401

RICE WEEVIL, (CALANDRA) SITOPHILUS ORYZA

Plate

60. *Sitophilus oryza:* A.—Egg. B.—Pupa, dorsal aspect. C.—Pupa, lateral aspect. D.—Pupa, ventral aspect. E.—Adult. F.—Third-stage larva. G.—First-stage larva. H.—Second-stage larva. I.—Fourth-stage larva. 422

OPIUS FLETCHERI AS A PARASITE OF THE MELON FLY IN HAWAII

Text Figures

1. *Opius fletcheri:* Egg just deposited 424
2. *Opius fletcheri:* Mature egg 424
3. *Opius fletcheri:* Larva, first instar, ventral aspect, showing head characters and complete tracheal system, and the egg serosal cells 425
4. *Opius fletcheri:* Molted skin of first-instar larva, showing the absence of egg serosal cells 426
5. *Opius fletcheri:* New second-instar larva 427
6. *Opius fletcheri:* Mandible of second-instar larva 427
7. *Opius fletcheri:* Mandible of third-instar larva 427
8. *Opius fletcheri:* Larva, fourth instar, lateral aspect, showing general outline and spiracles 428
9. *Opius fletcheri:* Spines on body of mature larva 428
10. *Opius fletcheri:* Mandible of fourth-instar larva 429
11. *Opius fletcheri:* Head of mature larva, dorso-cephalic aspect 429
12. *Opius fletcheri:* Pupa, female 430
13. *Opius fletcheri:* Adult female 431

TAMARIND POD-BORER, SITOPHILUS LINEARIS (HERBST)

Plate

61. *Sitophilus linearis:* A.—Pupa, dorsal view. B.—Pupa, front view. C.—Egg. D.—Mandible. E.—Mature larva. F.—Ventral view of head. G.—Clypeus and labrum. H.—Pupa, lateral view. I.—Head, face view. J.—Head, dorsal view. K.—Head, lateral view 446

INFLUENCE OF TEMPERATURE AND HUMIDITY ON THE GROWTH OF PSEUDOMONAS CITRI AND ITS HOST PLANTS AND ON INFECTION AND DEVELOPMENT OF THE DISEASE

Text Figure

1. Graph showing the rate of enzym action, as expressed in millimeters, at the various temperatures for a period of eight days on soluble starch agar... Page 451

DAUBENTONIA LONGIFOLIA (COFFEE BEAN), A POISONOUS PLANT

Plate

62. Herbarium specimen of *Daubentonia longifolia*, showing flowers, leaves, and pods..... 514

FUSARIUM-WILT OF TOBACCO

Text Figure

1. Camera-lucida drawings of spore forms of *Fusarium oxysporum* var. *nicotianae*, n. var.: A, macroconidia; B, microconidia; C, chlamydospores; D, conidiophore of the sporodochial stage..... 521

Plates

63. A.—A typical spot in a field of Maryland Broadleaf tobacco infested with Fusarium-wilt. Benedict, Md. 1916. B.—Uninoculated control. C.—Plants grown in soil artificially inoculated with the tobacco-wilt Fusarium and planted to White Burley..... 536

64. A.—Plant infected with Fusarium-wilt, showing wilting in vertical line on stalk. B.—Last stages of Fusarium-wilt in Maryland Broadleaf tobacco..... 536

65. A.—Result of plating out five pieces of infected vascular tissue from infected plant, illustrating character of growth of mycelium on potato agar. B.—Stem and midrib of plant, cut longitudinally to show the blackened vascular system..... 536

66. A.—Cross sections through vascular system of tobacco plant infected with Fusarium-wilt, showing the fungus mycelium in the vessels. B.—Longitudinal sections through the vascular system of plants infected with Fusarium-wilt, showing the fungus strands in the vessels..... 536

67. I.—Plants illustrating the influence of soil temperature on degree of wilting of plants in soil infested with Fusarium-wilt. II.—Plants grown in the same soil uninfested and at corresponding soil temperatures. III.—Plants illustrating the influence of varying soil reaction on the amount of Fusarium-wilt in infested soil. IV.—Plants illustrating varietal differences in resistance of tobacco to Fusarium-wilt..... 536

NODULE BACTERIA OF LEGUMINOUS PLANTS

Plates

68. A.—Soybean bacteria, J. K. Wilson's strain, 4 days old. B.—Vetch bacteria, 3 days old. C.—Bacillus radiobacter, 2 days old. D.—Soybean bacteria, beef agar, 4 days old. E.—Red clover bacteria, beef agar, 4 days old. F.—*Bacillus radiobacter*, beef agar, 4 days old. G.—Cowpea bacteria, potato, 6 days old. H.—Red clover bacteria, potato, 14 days old. I.—*B. radiobacter*, milk, 7 days old. J.—Cowpea bacteria, mannite-nitrate agar, 8 days old. K.—Vetch bacteria, mannite-nitrate agar, 8 days old. L.—*B. radiobacter*, mannite-nitrate solution, 17 days old.... 556

69. A.—Mannite-nitrate agar slants, 8 days old, from left to right: soybean bacteria, vetch bacteria, and *Bacillus radiobacter*. B.—Growth in milk, 4 weeks old, from left to right: soybean bacteria, vetch bacteria, and *B. radiobacter*. C.—Growth on potato, 2 weeks old: vetch bacteria (left) and *B. radiobacter* (right) ..... Page 556

CORRELATION AND CAUSATION

Text Figures

1. Diagram illustrating the interrelations among the factors which determine the weight of guinea pigs at birth and at weaning (33 days) 560
2. Diagram showing relations between two variables, X and Y, whose values are determined in part by common causes, B, C, and D, which are independent of each other 565
3. Diagram showing relations between two variables, X and Y, whose values are completely determined by common causes, B and C, which are independent of each other 565
4. A system in which the value of variable X is completely determined by causes M and N, which are correlated with each other 566
5. A system in which the value of X is affected by a factor, B, along two different paths, BMX and BNX 567
6. Diagram showing relations between two variables, X and Y, whose values are determined in part by common causes, M and N, which are correlated with each other 568
7. Simplified diagram of factors which determine birth weight in guinea pigs. 568
8. Path coefficients measuring the relations between birth rate (X), rate of growth (Q), gestation period (P), size of litter (L), and other causes (A, C) 570
9. Coefficients of determination. Symbols as in figure 7 570
10. Effect and one known cause 571
11. Effect and two correlated known causes 571
12. Effect and three correlated known causes 571
13. Effect and four correlated known causes 572
14. Relations between wet-bulb depression (B), wind velocity (W), radiation (R), and temperature (T) as assumed for direct analysis 576
15. Relations between factors of figure 14 and absolute humidity (H) expressing causal relations better than figure 14 but adapted only to indirect analysis. 579
16. Relations between evaporations or transpiration (X) and the system shown in figure 15 582

INHERITANCE OF SYNDACTYLISM, BLACK, AND DILUTION IN SWINE

Plate

70. The four types of F_2 segregates from a cross between mule-foot boar and Duroc-Jersey sows. A.—Black mule-foot. B.—Black cloven foot. C.—Red mule-foot. D.—Red cloven foot 604

FOUR RHYNCHOPHORA ATTACKING CORN IN STORAGE

Plates

71. *Araecerus fasciculatus*. A.—Pupa, dorsal view. B.—Pupa, front view. C.—Egg. D.—Mandible. E.—Mature larva. F.—Ventral view of head. G.—Labium and clypeus. H.—Pupa, lateral view. I.—Head, face view. J.—Head, dorsal view. K.—Head, lateral view 614

72. *Caulophilus latinasus:* A.—Pupa, dorsal view. B.—Pupa, front view. C.—Egg. D.—Mandible. E.—Mature larva. F.—Ventral view of head. G.—Labium and clypeus. H.—Pupa, lateral view. I.—Head, face view. J.—Head, dorsal view. K.—Head, lateral view..... Page 614

73. *Sitophilus oryza:* A.—Pupa, dorsal view. B.—Pupa, front view. C.—Egg. D.—Mandible. E.—Mature larva. F.—Ventral view of head. G.—Labium and clypeus. H.—Pupa, lateral view. I.—Head, face view. J.—Head, dorsal view. K.—Head, lateral view..... 614

74. *Sitophilus granarius:* A.—Pupa, dorsal view. B.—Pupa, front view. C.—Egg. D.—Mandible. E.—Mature larva. F.—Ventral view of head. G.—Labium and clypeus. H.—Pupa, lateral view. I.—Head, face view. J.—Head, dorsal view. K.—Head, lateral view..... 614

FREEZING OF FRUIT BUDS

Plate

75. Apparatus for freezing entire tree..... 662

EFFECT OF VARIOUS CROPS UPON THE WATER EXTRACT OF A TYPICAL SILTY CLAY LOAM SOIL

Text Figures

1. Decrease of water-soluble nutrients from the growth of various crops, as shown by increases in specific resistance..... 664
2. Decrease of water-soluble nutrients from varying numbers of barley plants, as shown by increase in specific resistance..... 664
3. Decrease of water-soluble nitrates from the growth of various crops. (Graphs $=\frac{1}{2}$ NO_3.)..... 665
4. Decrease of water-soluble nitrates from varying numbers of barley plants. (Graphs$=\frac{1}{2}$ NO_3.)..... 666
5. Decrease in the concentration of soil solution shown by freezing-point depression..... 666

ANOTHER CONIDIAL SCLEROSPORA OF PHILIPPINE MAIZE

Text Figure

1. Comparison of the sizes of 700 conidia of *Sclerospora spontanea* with 700 conidia of *S. philippinensis; A*, variation of conidia in length; *B*, variation of conidia in diameter; *C*, ratios of length to width of conidia arranged in classes..... 675

Plates

76. Corner of a native-grown maize plot in the interior uplands of Cebu..... 684
77. A.—Clump of *Saccharum spontaneum*, showing characteristic size and habit of healthy plants under natural conditions. B.—Clump of *Saccharum spontaneum* infected with *Sclerospora spontanea*..... 684
78. A.—A young seedling (3 weeks old) of *Saccharum spontaneum* infected with *Sclerospora spontanea*. B.—Conidiophores on the leaf of *Saccharum spontaneum*. C.—Young shoots of *Saccharum spontaneum* arising after the primary stalk had been cut, and like it severely infected with *Sclerospora spontanea*..... 684

79. A.—Typical conidiophore, showing characteristically long, slender, unknobbed basal cell, relatively short main axis with its greatest diameter about midway to the primary branches, and fairly well-developed branch system bearing long, slender conidia. B.—Upper portion of a conidiophore which has a poorly developed branch system and hence bears few conidia on sterigmata which are relatively large. C.—Portion of the branch system of a conidiophore, showing the conidia germinating while still attached to their sterigmata. D.—Stalk portion of a typical conidiophore, showing long, slender, unknobbed basal cell, and main axis which is slender above the septum, expands rapidly to its greatest diameter about midway, and contracts again below the branches. E, F.—Typical basal cells of conidiophores. G.—Stalk portion of a conidiophore with basal cell which, though unusually short, nevertheless is longer than the extent of the main axis from septum to primary branches. H.—Typical stalk portion of a conidiophore from sugar cane. I, J, K.—Typical conidia showing variations in size and shape and method of germination by hyphae ..... Page 684

ONION SMUDGE

Text Figures

1. Conidia and appressoria of *Colletotrichum circinans* 689
2. Acervulus of *Colletotrichum circinans* on artificially inoculated onion scale. 690
3. Spores of *Colletotrichum fructus* (*A*) and *C. circinans* (*B*) 694
4. Relation of temperature to growth of *Colletotrichum circinans* on agar plates. 697
5. Relation of temperature to spore germination of *Colletotrichum circinans* ... 698
6. *Colletotrichum circinans:* Stage of penetration of epidermal cell of onion scale at 66 hours after inoculation 702
7. Cross section of epidermis, showing early stage of penetration by *Colletotrichum circinans* 703
8. Cross section of epidermis (*A*) and underlying parenchyma cells (*B*) of onion scale inoculated with a suspension of *Colletotrichum circinans* spores and kept in a moist chamber at room temperature 704
9. Cross section of onion scale naturally infected with *Colletotrichum circinans*, showing the mycelium developing first just beneath the cuticle and later penetrating the subcuticular wall 705
10. Chart from data collected at Racine, Wis., during 1915 and 1916, showing the daily mean soil temperature at a depth of 1 to 2 inches, and the rainfall. 708

Plates

80. Onion smudge: Onion sets (White Portugal variety) naturally infected with *Colletotrichum circinans* 722
81. Onion smudge: A, B, E, D.—Advanced stages of smudge after several months in storage. C.—Bulb inoculated in a moist chamber with a suspension of *Colletotrichum circinans* conidia. F, G.—*Macrosporium* sp. on outer scale of white onion sets. H.—*M. porrum* and *Phoma alliicola* on outer scale of white onion set 722
82. Relation of soil temperature to the development of smudge 722
83. *Colletotrichum circinans* and *C. fructus:* A.—Photomicrograph of cross section of naturally infected onion scale. B.—Photomicrograph of cross section of an infected onion scale held for several months in poorly ventilated storage. C, D.—Photomicrographs of cross sections of *C. circinans* (C) and *C. frutus* (D) on apple fruit 722

84. *Colletotrichum fructus* and *C. circinans:* A.—Dilution plate from spores of *Colletotrichum fructus*. B.—Individual colony of *C. fructus* on potato agar. C.—Apple of Fameuse variety inoculated with mycelium from pure culture of *C. circinans*. D.—Dilution plate from spores of *C. circinans*. E.—Individual colony of *C. circinans* on potato agar..... Page 722

85. Relation of curing conditions to the development of smudge: A, B.—Comparison of onion sets artificially dried immediately after harvest with those not dried. C, D.—Comparison of white onion sets cured in shallow crates in the field under the best of natural conditions with part of the same lot after exposure to moist conditions for one week..... 722

VARIATIONS IN COLLETOTRICHUM GLOEOSPORIOIDES

Plate

86. A, B.—Variation occurring in strain 990. The cultures were not made from a single spore. C.—Variation occurring in a culture of strain 990 which was made from a single spore..... 736

Text Figures

1. Variability of strains of *Colletotrichum gloeosporioides* in spore length..... 728

2. I, culture 510: A, greenish black mycelium; B, white mycelium. II, culture 943: A, black mycelium; B, white mycelium; C, mycelium mostly in medium, growth zoned, abundant spore production. III, culture 495: A, black mycelium; B, gray mycelium; C, white mycelium. IV, culture 527: A, gray mycelium; B, greenish black mycelium; C, white mycelium; D, black mycelium. V, culture 940: A, greenish black mycelium; B, white mycelium, some greenish concentric circles; C, black mycelium; D, white mycelium; E, white and black mixed..... 734

A TRANSMISSIBLE MOSAIC DISEASE OF LETTUCE

Plate

87. A.—Leaves of Romaine lettuce. B.—Young expanding leaves of head lettuce from experiment started March 22..... 740

LECONTE'S SAWFLY, AN ENEMY OF YOUNG PINES

Text Figures

1. Chart showing life and seasonal history of *Neodiprion lecontei* through the active period of three years (November to March omitted, the insect being in the cocoon during this period)..... 751

2. Position of end of abdomen of female when ovipositing, showing the various parts and their position: 1, lance; 2, apical part of sheath; 3, basal part of sheath; 4, nates or ninth tergite; 5, eighth sternite; 6, chitinized rods at base of lancet; 7, lancet..... 755

3. Distribution of *Neodiprion lecontei*. The larger dots indicate places from which specimens have actually been received..... 759

Plates

88. *Neodiprion lecontei:* A.—Adult female. B.—Adult male..... 760

89. *Neodiprion lecontei:* A.—Larva. B.—Sixth-stage larva: The muscles of a single abdominal segment distributed over several segments to show their numbers, position, and attachment..... 760

90. *Neodiprion lecontei:* Sixth-stage larva. A.—Front view of head. B.—Ventral (or apical) view of head capsule. C.—Front view of head capsule. D.—Lateral view of head. E.—Sagittal section of head. F.—Antenna. G.—Frons, adfrons, and clypeus. H.—Mandibles. I.—Epipharynx and labrum. J.—Internal view of hypopharynx maxillæ, and labium. K.—External view of maxillæ, and labium. L.—External view of maxillæ. M.—Interior and apical view of maxilla. N.— Page
End view of maxilla. O.—End view of labium..... 760
91. *Neodiprion lecontei:* Sixth-stage larva. A.—External view of the thorax. B.—External view of the second and third abdominal segments. C.—External view of the ninth and tenth abdominal segments. D.—Internal view of thoracic skin. E.—Internal view of the skin of the second and third abdominal segments. F.—Diagrammatic cross section of the abdomen showing the longitudinal areas of the body on its transverse circumference..... 760
92. *Neodiprion lecontei:* A.—Some defoliated twigs showing feeding on bark of stem. B.—Eggs within needles of *Pinus virginiana*..... 760

EVALUATION OF CLIMATIC TEMPERATURE EFFICIENCY FOR THE RIPENING PROCESSES IN SWEETCORN

Text Figure

1. Comparison of early and late crops of sweet corn in respect to changes in percentage composition in equal lengths of time..... 798

SOME LEPIDOPTERA LIKELY TO BE CONFUSED WITH THE PINK BOLLWORM

Plates

93. Male genitalia (Gelechiidae): A.—*Gelechia trophella:* Posterior part of tegumen, showing uncus and gnathos, ventral view. B.—*G. trophella:* Lateral view of male genitalia with eighth abdominal segment attached. C.—*G. hibiscella:* Lateral view of male genitalia with eighth abodminal segment attached..... 836
94. Male genitalia (Gelechiidae): A.—*Telphusa mariona* (type): Lateral view of male genitalia. B.—*T. mariona* (type): Posterior part of tegumen, showing uncus, ventral view. C.—*Gelechia neotrophella* (type): Aedoeagus and penis. D.—*G. neotrophella* (type): Lateral view of male genitalia with aedoeagus and eighth segment removed. E.—*G. neotrophella* (type): Posterior part of tegumen, showing uncus and gnathos, ventral view. F.—*G. neotrophella* (type): Posterior half of harpes, ventral view. G.—*G. neotrophella* (type): Sternite and tergite of modified eighth abdominal segment..... 836
95. Male genitalia (Gelechiidae, Stenomidae, and Oecophoridae): A.—*Isophrictis similiella:* Ventral view of male genitalia, spread. B.—*Aedemoses haesitans:* Ventral view of male genitalia, spread. C.—*A. haesitans:* Enlargement of typical split hair on cucullus. D.—*Borkhausenia fasciata:* Ventro-lateral view of male genitalia, spread, showing a symmetrical armlike projections from gnathos and costa of harpes..... 836
96. Male genitalla (Oecophoridae): A.—*Borkhausenia minutella:* Aedoeagus. B.—*B. minutella:* Ventral view of male genitalia, spread, aedoeagus omitted. C.—*B. diveni* (type): Ventral view of male genitalia, spread. D.—*B. diveni* (type): Dorsal view of an abdominal segment showing spinose condition of abdomen. E.—*B. diveni* (type): Modified tergite of eighth abdominal segment. F.—*B. diveni* (type): Modified sternite of eighth abdominal segment..... 836

97. Male genitalia (Oecophoridae): A.—*Borkhausenia conia:* Portion of tergite of seventh abdominal segment, showing spinose and chitinized character of caudal margin. B.—*B. conia:* Ventral view of male genitalia, spread, aedoeagus omitted. C.—*B. conia:* Aedoeagus. D.—*B. conia:* Modified tergite of eighth abdominal segment. E.—*B. conia:* Modified sternite of eighth abdominal segment..... Page 836

98. Malegenitalia (Blastobasidae): A.—*Zenodochium citricolella:* Aedoeagus. B.—*Z. citricolella:* Lateral view of male genitalia, right harpe and aedoeagus omitted. C.—*Z. citricolella:* Right harpe. D.—*Holcocera ochrocephala:* Ventral view of male genitalia, spread, aedoeagus omitted. E.—*H. ochrocephala:* Dorsum of an abdominal segment showing transverse row of spines. F.—*H. ochrocephala:* Aedoeagus and penis..... 836

99. Male genitalia (Olethreutidae and Blastobasidae): A.—*Crocidosema plebeiana:* Ventral view of male genitalia, spread. B.—*Eucosma discretivana* (type): Ventral view of male genitalia, spread. C.—*Holcocera confamulella* (type): Ventral view of male genitalia, spread..... 836

100. Male genitalia (Phaloniidae and Pryalidae): A.—*Phalonia cephalanthana* (type): Ventral view of male genitalia, spread. B.—*Homoeosoma electellum:* Ventral view of male genitalia, spread..... 836

101. Larval structures: A.—*Pectinophora gossypiella:* Head capsule, dorsal view, showing arrangement of setæ. B.—*P. gossypiella:* Head capsule, lateral view, showing arrangement of setæ. C.—*Dicymolomia julianalis:* Head capsule, dorsal view, showing arrangement of setæ. D.—*D. julianalis:* Head capsule, lateral view, showing arrangement of setæ. E.—*Meskea dyspteraria:* Head capsule, dorsal view, showing arrangement of setæ. F.—*M. dyspteraria:* Head capsule, lateral view, showing arrangement of setæ..... 836

102. Larval structures: A.—*Pyroderces rileyi:* Head capsule, dorsal view, showing arrangement of setæ. B.—*P. rileyi:* Head capsule, lateral view, showing arrangement of setæ. C.—*Crocidosema plebeiana:* Head capsule, dorsal view, showing arrangement of setæ. D.—*C. plebeiana:* Head capsule, lateral view, showing arrangement of setæ. E..—*Z. enodochium citricolella:* Labium and maxillæ. F.—*Isophrictis similiella:* Head capsule, dorsal view, showing arrangement of setæ..... 836

103. Larval structures: A.—*Pectinophora gossypiella:* Setal maps of first and second thoracic and third, eighth, and ninth abdominal segments. B.—*Dicymolomia julianalis:* Setal maps of first and second thoracic and third, eighth, and ninth abdominal segments. C.—*Pyroderces rileyi:* Setal maps of first thoracic and eighth and ninth abdominal segments. D.—*Heliothis obsoleta:* Setal maps of first thoracic and third abdominal segments. E.—*Crocidosema plebeiana:* Setal maps of first and second thoracic and third, eighth, and ninth abdominal segments..... 836

104. Larval structures: A.—*Platynota rostrana:* Setal maps of first and second thoracic and third, eighth, and ninth abdominal segments. B.—*Meskea dyspteraria:* Setal maps of first and second thoracic and third, eighth, and ninth abdominal segments. C.—*Z. enodochium citricolella:* Setal maps of first thoracic and third, eighth, and ninth abdominal segments. D.—*Aedemoses haesitans:* Setal map of third abdominal segment. E.—*Moodna estrinella:* Setal maps of second thoracic and eighth and ninth abdominal segments..... 836

105. Larval structures: A.—*Platynota rostrana:* Setal maps of eighth and ninth abdominal segments, dorsal view. B.—*Eucosma helianthana:* Setal maps of eighth and ninth abdominal segments, dorsal view. C.—*Pectinophora gossypiella:* Setal maps of eighth and ninth abdominal segments, dorsal view. D.—*Pyroderces rileyi:* Setal maps of eighth snd ninth abdominal segments, dorsal view. E.—*Pectinophora gossypiella:* Prothorax, ventral view, showing position of legs. F.—*Telphusa mariona:* Ventro-caudal view of tenth abdominal segment, showing anal fork. G.—*Crocidosema plebeiana:* Ventro-caudal view of tenth abdominal segment, showing anal fork. H.—*Gelechia neotrophella:* Ventro-caudal view of tenth abdominal segment, showing anal fork. I.—*Zenodochium citricolella:* Prothorax, ventral view, showing position of legs. Page 836

106. Larval structures: A.—*Pectinophora gossypiella:* Crochet arrangement of abdominal prolegs. B.—*Crocidosema plebeiana:* Crochet arrangement of abdominal prolegs. C.—*Pyroderces rileyi:* Crochet arrangement of abdominal prolegs. D.—*Dicymolomia julianalis:* Crochet arrangement of abdominal proleg. E.—*Heliothis obsoleta:* Crochet arrangement of abdominal proleg. 836

107. Pupal structures: A.—*Pectinophora gossypiella:* Ventral view of pupa. B.—*Pectinophora gossypiella:* Caudal end of pupa, lateral view. C.—*Pectinophora gossypiella:* Mature pupa, ventral view, shaded to show eyes of imago visible through pupal skin and characteristic pubescence of the pupa. D.—*Pectinophora gossypiella:* Dorsal view of pupa. E.—*Pyroderces rileyi:* Ventral view of pupa. F.—*Pyroderces rileyi:* Dorsal view of pupa. 836

108. Pupal structures: A.—*Crocidosema plebeiana:* Abdomen of female pupa, ventral view. B.—*C. plebeiana:* Abdomen of male pupa, ventral view. C.—*C. plebeiana:* Lateral view of an abdominal segment, showing arrangement and character of dorsal spines; one spine greatly enlarged to show shape. D.—*C. plebeiana:* Abdomen of pupa, dorsal view. E.—*Dicymolomia julianalis:* Dorsal view of pupa. F.—*D. julianalis:* Caudal end of pupa, lateral view. G.—*D. julianalis:* Caudal end of male pupa, ventral view. H. *D. julianalis:* Ventral view of female pupa. 836

109. Pupal structures: A.—*Meskea dyspteraria:* Caudal end of female pupa, lateral view. B.—*M. dyspteraria:* Abdomen of female pupa, ventral view. C.—*M. dyspteraria:* Male pupa, dorsal view. D.—*M. dyspteraria:* Caudal end of male pupa, lateral view. E.—*M. dyspteraria:* Male pupa, ventral view. F.—*Amorbia emigratella:* Abdomen of pupa, dorsal view. G.—*Telphusa mariona:* Caudal end of pupa, ventral view, showing peculiarly scalloped and fringed caudal margin of seventh abdominal segment. 836